

EXTRA INVARIANCE OF SHIFT-INVARIANT SPACES ON LCA GROUPS

M. ANASTASIO, C. CABRELLI, AND V. PATERNOSTRO

ABSTRACT. Let G be an LCA group and K a closed subgroup of G . A closed subspace of $L^2(G)$ is called K -invariant if it is invariant under translations by elements of K .

Assume now that H is a countable uniform lattice in G and M is any closed subgroup of G containing H . In this article we study necessary and sufficient conditions for an H -invariant space to be M -invariant.

As a consequence of our results we prove that for each closed subgroup M of G containing the lattice H , there exists an H -invariant space S that is exactly M -invariant. That is, S is not invariant under any other subgroup M' containing M . We also obtain estimates on the support of the Fourier transform of the generators of the H -invariant space, related to its M -invariance.

1. INTRODUCTION

Let G be an LCA group and $K \subseteq G$ a subgroup. For $y \in G$ let us denote by t_y the translation operator acting on $L^2(G)$. That is, $t_y f(x) = f(x - y)$ for $x \in G$ and $f \in L^2(G)$.

A closed subspace S of $L^2(G)$ satisfying that $t_k f \in S$ for every $f \in S$ and every $k \in K$ is called K -invariant.

In case that G is \mathbb{R}^d and K is \mathbb{Z}^d the subspace S is called shift-invariant. Shift-invariant spaces are central in several areas such as approximation theory, wavelets, frames and sampling.

The structure of these spaces for the group \mathbb{R}^d has been studied in [Hel64, Sri64, HS64, dBDR94a, dBDR94b, Bow00] and, in the context of general LCA groups, in [KR08, CP09].

Let now $H \subseteq G$ be a countable uniform lattice and M be any closed subgroup of G satisfying that $H \subseteq M \subseteq G$. In this article we want to study necessary and sufficient conditions in order that an H -invariant space $S \subseteq L^2(G)$ is M -invariant.

For the case $(\mathbb{Z}, M, \mathbb{R})$, this has been studied by Aldroubi et al in [ACHKM09] and for the case $(\mathbb{Z}^d, M, \mathbb{R}^d)$ by Anastasio et al in [ACP09].

Date: June 5, 2009.

2000 Mathematics Subject Classification. Primary 43A77; Secondary 43A15.

Key words and phrases. Shift-invariant space, translation invariant space, LCA groups, range functions, fibers.

The research of the authors is partially supported by Grants: ANPCyT, PICT 2006–177, CONICET, PIP 5650, UBACyT X058 and X108.

In this article we consider the extension to general LCA groups. This is important in order to obtain general conditions that can be applied to different cases, as for example the case of the classic groups as the d -dimensional torus \mathbb{T}^d , the discrete group \mathbb{Z}^d , and the finite group \mathbb{Z}_d .

This article is organized as follows. In Section 2 we set the notation and give some definitions. We study the properties of the invariance set in Section 3. In Section 4 we describe the structure of M -invariant spaces and range functions in the context of LCA groups. The characterizations of M -invariance for H -invariant spaces is given in section 5. Finally in Section 6 we give some applications.

2. NOTATION

Let G be an arbitrary locally compact Hausdorff abelian group (LCA) written additively. We will denote by m_G its Haar measure. The dual group of G , that is, the set of continuous characters on G , is denoted by \widehat{G} . The value of the character $\gamma \in \widehat{G}$ at the point $x \in G$, is written by (x, γ) .

The Fourier transform of a Haar integrable function f on G , is the function \widehat{f} on \widehat{G} defined by

$$\widehat{f}(\gamma) = \int_G f(x)(x, -\gamma) dm_G(x), \quad \gamma \in \widehat{G}.$$

When the Haar measures m_G and $m_{\widehat{G}}$ are normalized such that the Inversion Formula holds (see [Rud62]), the Fourier transform on $L^1(G) \cap L^2(G)$ can be extended to a unitary operator from $L^2(G)$ onto $L^2(\widehat{G})$, the so-called Plancharel transformation. We also denote this transformation by “ \wedge ”.

Note that the Fourier transform satisfies $t_x \widehat{f}(\cdot) = (-x, \cdot) \widehat{f}(\cdot)$.

For a subgroup K of G , the set

$$K^* = \{\gamma \in \widehat{G} : (k, \gamma) = 1, \forall k \in K\}$$

is called the *annihilator* of K . Since every character in \widehat{G} is continuous, K^* is a closed subgroup of \widehat{G} .

We will say that a closed subspace $V \subseteq L^2(G)$ is *K -invariant* if

$$f \in V \Rightarrow t_k f \in V \quad \forall k \in K.$$

For a subset $\mathcal{A} \subseteq L^2(G)$, define

$$E_K(\mathcal{A}) = \{t_k \varphi : \varphi \in \mathcal{A}, k \in K\} \quad \text{and} \quad S_K(\mathcal{A}) = \overline{\text{span}} E_K(\mathcal{A}).$$

We call $S_K(\mathcal{A})$ the K -invariant space generated by \mathcal{A} . If $\mathcal{A} = \{\varphi\}$, we simply write $S_K(\varphi)$, and we call $S_K(\varphi)$ a *principal K -invariant space*.

Let L be a subset of G , we will say that a function f defined on G is L -periodic if $t_\ell f = f$ for all $\ell \in L$. A subset $B \subseteq G$ is L -periodic if its indicator function (denoted by χ_B) is L -periodic.

When two LCA groups G_1 and G_2 are topologically isomorphic we will write $G_1 \approx G_2$.

3. THE INVARIANCE SET

Here and subsequently G will be an LCA group and H a countable uniform lattice in G . That is, a countable discrete subgroup of G with compact quotient group G/H .

For simplicity of notation throughout this paper we will write Γ instead of \widehat{G} .

The aim of this work is to characterize the extra invariance of an H -invariant space. For this, given $S \subseteq L^2(G)$ an H -invariant space, we define the *invariance set* as

$$M = \{x \in G : t_x f \in S, \forall f \in S\}. \quad (1)$$

If \mathcal{A} is a set of generators for S , it is easy to check that $m \in M$ if and only if $t_m \varphi \in S$ for all $\varphi \in \mathcal{A}$.

In case that $M = G$, Wiener's theorem (see [Hel64], [Sri64] and [HS64]) states that there exists a measurable set $E \subseteq \Gamma$ satisfying

$$S = \{f \in L^2(G) : \text{supp}(\widehat{f}) \subseteq E\}.$$

We want to describe S when M is not all G . We will first study the structure of the set M .

Proposition 3.1. *Let S be an H -invariant space of $L^2(G)$ and let M be defined as in (1). Then M is a closed subgroup of G containing H .*

For the proof of this proposition we will need the following lemma. Recall that a semigroup is a nonempty set with an associative additive operation.

Lemma 3.2. *Let K be a closed semigroup of G containing H , then K is a group.*

Proof. Let π be the quotient map from G onto G/H . Since K is a semigroup containing H , we have that $K + H = K$, thus

$$\pi^{-1}(\pi(K)) = \bigcup_{k \in K} k + H = K + H = K. \quad (2)$$

This shows that $\pi(K)$ is closed in G/H and therefore compact.

By [HR63, Theorem 9.16], we have that a compact semigroup of G/H is necessarily a group, thus $\pi(K)$ is a group and consequently K is a group. \square

Proof of Proposition 3.1. Since S is an H -invariant space, $H \subseteq M$.

We first proceed to show that M is closed. Let $x_0 \in G$ and $\{x_\lambda\}_{\lambda \in \Lambda}$ a net in M converging to x_0 . Then

$$\lim_{\lambda} \|t_{x_\lambda} f - t_{x_0} f\|_2 = 0.$$

Since S is closed, it follows that $t_{x_0} f \in S$, thus $x_0 \in M$.

It is easy to check that M is a semigroup of G , hence we conclude from Lemma 3.2 that M is a group. \square

4. THE STRUCTURE OF PRINCIPAL M -INVARIANT SPACES

4.1. Preliminaries. Shift-invariant spaces in $L^2(\mathbb{R}^d)$ are completely characterized using fiberization techniques and range functions (see [Bow00]). This theory has been extended to general LCA groups in [CP09]. In what follows we state some definitions and properties given in that work.

We will assume that G is a second countable LCA group and H a countable uniform lattice in G .

The fact that G is second countable, G/H is compact and $\widehat{G/H} \approx H^*$, implies that H^* is countable and discrete. Moreover, since $\Gamma/H^* \approx \widehat{H}$, H^* is a countable uniform lattice in Γ . Therefore, there exists a measurable section Ω of Γ/H^* with finite m_Γ -measure (see [KK98] and [FG64]).

Let $L^2(\Omega, \ell^2(H^*))$ be the space of all measurable functions $\Phi : \Omega \rightarrow \ell^2(H^*)$ such that

$$\|\Phi\|_2^2 := \int_{\Omega} \|\Phi(\omega)\|_{\ell^2(H^*)}^2 dm_\Gamma(\omega) < \infty.$$

The following proposition shows that the space $L^2(\Omega, \ell^2(H^*))$ is isometric (up to a constant) to $L^2(G)$.

Proposition 4.1. *The mapping $\mathcal{T}_H : L^2(G) \rightarrow L^2(\Omega, \ell^2(H^*))$ defined as*

$$\mathcal{T}_H f(\omega) = \{\widehat{f}(\omega + h^*)\}_{h^* \in H^*}$$

is an isomorphism that satisfies $\|\mathcal{T}_H f\|_2 = m_{H^}(\{0\})^{1/2} \|f\|_{L^2(G)}$.*

For $f \in L^2(G)$, the sequence $\mathcal{T}_H f(\omega) = \{\widehat{f}(\omega + h^*)\}_{h^* \in H^*}$ is the H -fiber of f at ω . Given a subspace V of $L^2(G)$ and $\omega \in \Omega$, the H -fiber space of V at ω is

$$J_H(V)(\omega) = \overline{\{\mathcal{T}_H f(\omega) : f \in V\}},$$

where the closure is taken in the norm of $\ell^2(H^*)$.

The map that assigns to each ω the fiber space $J_H(V)(\omega)$ is known in the literature as the *range function* of V .

The following proposition characterizes H -invariant spaces in terms of range functions and fibers.

Proposition 4.2. *If S is an H -invariant space in $L^2(G)$, then*

$$S = \{f \in L^2(G) : \mathcal{T}_H f(\omega) \in J_H(S)(\omega) \text{ for a.e. } \omega \in \Omega\}.$$

Moreover, if $S = S_H(\mathcal{A})$ for a countable set $\mathcal{A} \subseteq L^2(G)$, then, for almost every $\omega \in \Omega$,

$$J_H(S)(\omega) = \overline{\text{span}}\{\mathcal{T}_H \varphi(\omega) : \varphi \in \mathcal{A}\}.$$

In this article we will use fiberizations techniques for a more general case, since the subspaces will be invariant under a closed subgroup which is not necessarily discrete.

The above results from [CP09] can be extended straightforward to the case that the spaces are invariant under a closed subgroup M of G containing a countable uniform lattice H as follows.

Since $H \subseteq M$, we have that $M^* \subseteq H^*$ and, in particular, M^* is discrete. Thus, there exists a countable section \mathcal{N} of H^*/M^* . Then, the set given by

$$\mathcal{D} = \bigcup_{\sigma \in \mathcal{N}} \Omega + \sigma \quad (3)$$

is a σ -finite measurable section of the quotient Γ/M^* . Using this section of Γ/M^* it is possible to obtain, in a way analogous to the discrete case, the following.

Proposition 4.3.

i) The mapping $\mathcal{T}_M : L^2(G) \longrightarrow L^2(\mathcal{D}, \ell^2(M^*))$ defined as

$$\mathcal{T}_M f(\delta) = \{\widehat{f}(\delta + m^*)\}_{m^* \in M^*}$$

is an isomorphism that satisfies $\|\mathcal{T}_M f\|_2 = m_{M^*}(\{0\})^{1/2} \|f\|_{L^2(G)}$.

ii) Let S be an M -invariant space generated by a countable set \mathcal{A} . For each $\delta \in \mathcal{D}$, define the M -fiber space of S at δ as

$$J_M(S)(\delta) = \overline{\text{span}}\{\mathcal{T}_M \varphi(\delta) : \varphi \in \mathcal{A}\}.$$

If P and P_δ are the orthogonal projections onto S and $J_M(S)(\delta)$ respectively, then, for every $g \in L^2(G)$,

$$\mathcal{T}_M(Pg)(\delta) = P_\delta(\mathcal{T}_M g(\delta)) \quad \text{a.e. } \delta \in \mathcal{D}.$$

iii) If S is an M -invariant space in $L^2(G)$, then

$$S = \{f \in L^2(G) : \mathcal{T}_M f(\delta) \in J_M(S)(\delta) \text{ for a.e. } \delta \in \mathcal{D}\}.$$

4.2. Principal M -invariant Spaces.

We prove now the following characterization of principal M -invariant spaces. This result extends the \mathbb{R}^d case.

Theorem 4.4. Let $f \in L^2(G)$ and M a closed subgroup of G containing H . If $g \in S_M(f)$, then there exists an M^* -periodic function η such that $\widehat{g} = \eta \widehat{f}$.

Conversely, if η is an M^* -periodic function such that $\eta \widehat{f} \in L^2(\Gamma)$, then the function g defined by $\widehat{g} = \eta \widehat{f}$ belongs to $S_M(f)$.

Proof. Let us call $S = S_M(f)$ and let P and P_δ be the orthogonal projections onto S and $J_M(S)(\delta)$ respectively. Given $g \in S$, we first define η_g in \mathcal{D} as

$$\eta_g(\delta) = \begin{cases} \frac{\langle \mathcal{T}_M g(\delta), \mathcal{T}_M f(\delta) \rangle}{\|\mathcal{T}_M f(\delta)\|_2^2} & \text{if } \delta \in E_f \\ 0 & \text{otherwise,} \end{cases}$$

where E_f is the set $\{\delta \in \mathcal{D} : \|\mathcal{T}_M f(\delta)\|_2^2 \neq 0\}$. Then, since $\{\mathcal{D} + m^*\}_{m^* \in M^*}$ forms a partition of Γ , we can extend η_g to all Γ in an M^* -periodic way.

Now, by Proposition 4.3 we have that

$$\mathcal{T}_M g(\delta) = \mathcal{T}_M(Pg)(\delta) = P_\delta(\mathcal{T}_M g(\delta)) = \eta_g(\delta) \mathcal{T}_M f(\delta).$$

Since η_g is an M^* -periodic function, $\widehat{g} = \eta_g \widehat{f}$ as we wanted to prove.

Conversely, if $\widehat{g} = \eta \widehat{f}$, with η an M^* -periodic function, then $\mathcal{T}_M g(\delta) = \eta(\delta) \mathcal{T}_M f(\delta)$. By Proposition 4.3, $g \in S$. \square

5. CHARACTERIZATION OF M -INVARIANCE

If $H \subseteq M \subseteq G$, where H is a countable uniform lattice in G and M is a closed subgroup of G , we are interested in describing when an H -invariant space S is also M -invariant.

Let Ω be a measurable section of Γ/H^* and \mathcal{N} a countable section of H^*/M^* . For $\sigma \in \mathcal{N}$ we define the set B_σ as

$$B_\sigma = \Omega + \sigma + M^* = \bigcup_{m^* \in M^*} (\Omega + \sigma) + m^*. \quad (4)$$

Therefore, each B_σ is an M^* -periodic set.

Since Ω tiles Γ by H^* translations and \mathcal{N} tiles H^* by M^* translations, it follows that $\{B_\sigma\}_{\sigma \in \mathcal{N}}$ is a partition of Γ .

Now, given an H -invariant space S , for each $\sigma \in \mathcal{N}$, we define the subspaces

$$U_\sigma = \{f \in L^2(G) : \widehat{f} = \chi_{B_\sigma} \widehat{g}, \text{ with } g \in S\}. \quad (5)$$

5.1. Characterization of M -invariance in terms of subspaces.

The main theorem of this section characterizes the M -invariance of S in terms of the subspaces U_σ .

Theorem 5.1. *If $S \subseteq L^2(G)$ is an H -invariant space and M is a closed subgroup of G containing H , then the following are equivalent.*

- i) S is M -invariant.
- ii) $U_\sigma \subseteq S$ for all $\sigma \in \mathcal{N}$.

Moreover, in case any of these hold we have that S is the orthogonal direct sum

$$S = \bigoplus_{\sigma \in \mathcal{N}} U_\sigma.$$

Now we state a lemma that we need to prove Theorem 5.1.

Lemma 5.2. *Let S be an H -invariant space and $\sigma \in \mathcal{N}$. Assume that the subspace U_σ defined in (5) satisfies $U_\sigma \subseteq S$. Then, U_σ is an M -invariant space and in particular is H -invariant.*

Proof. Let us prove first that U_σ is closed. Suppose that $f_j \in U_\sigma$ and $f_j \rightarrow f$ in $L^2(G)$. Since $U_\sigma \subseteq S$ and S is closed, f must be in S . Further,

$$\|\widehat{f}_j - \widehat{f}\|_2^2 = \|(\widehat{f}_j - \widehat{f})\chi_{B_\sigma}\|_2^2 + \|(\widehat{f}_j - \widehat{f})\chi_{B_\sigma^c}\|_2^2 = \|\widehat{f}_j - \widehat{f}\chi_{B_\sigma}\|_2^2 + \|\widehat{f}\chi_{B_\sigma^c}\|_2^2.$$

Since the left-hand side converges to zero, we must have that $\widehat{f}\chi_{B_\sigma^c} = 0$ a.e. $\gamma \in \Gamma$. Then, $\widehat{f} = \widehat{f}\chi_{B_\sigma}$. Consequently $f \in U_\sigma$, so U_σ is closed.

Now we show that U_σ is M -invariant. Given $m \in M$ and $f \in U_\sigma$, we will prove that $(m, \cdot)\widehat{f}(\cdot) \in \widehat{U}_\sigma$.

Since $f \in U_\sigma$, there exists $g \in S$ such that $\widehat{f} = \chi_{B_\sigma} \widehat{g}$. Hence,

$$(m, \cdot)\widehat{f}(\cdot) = (m, \cdot)(\chi_{B_\sigma} \widehat{g})(\cdot) = \chi_{B_\sigma}(\cdot)((m, \cdot)\widehat{g}(\cdot)). \quad (6)$$

If we were able to find an H^* -periodic function ℓ_m verifying

$$(m, \gamma) = \ell_m(\gamma) \quad \text{a.e. } \gamma \in B_\sigma, \quad (7)$$

then, we can rewrite (6) as

$$(m, \cdot) \widehat{f}(\cdot) = \chi_{B_\sigma}(\cdot) (\ell_m \widehat{g})(\cdot).$$

Theorem 4.4 can then be applied for the uniform lattice H . Thus, since ℓ_m is H^* -periodic, we obtain that $\ell_m \widehat{g} \in \widehat{S_H(g)} \subseteq \widehat{S}$ and so, $(m, \cdot) \widehat{f}(\cdot) \in \widehat{U}_\sigma$.

Now we define the function ℓ_m as follows. For each $h^* \in H^*$, set

$$\ell_m(\omega + h^*) = (m, \omega + \sigma) \quad \text{a.e. } \omega \in \Omega. \quad (8)$$

It is clear that ℓ_m is H^* -periodic.

Since (m, \cdot) is M^* -periodic,

$$(m, \omega + \sigma) = (m, \omega + \sigma + m^*) \quad \text{a.e. } \omega \in \Omega, \forall m^* \in M^*.$$

Thus, (7) holds.

Note that, since $H \subseteq M$, the H -invariance of U_σ is a consequence of the M -invariance. □

Proof of Theorem 5.1. i) \Rightarrow ii): Fix $\sigma \in \mathcal{N}$ and $f \in U_\sigma$. Then $\widehat{f} = \chi_{B_\sigma} \widehat{g}$ for some $g \in S$. Since χ_{B_σ} is an M^* -periodic function, by Theorem 4.4, we have that $f \in S_M(g) \subseteq S$, as we wanted to prove.

ii) \Rightarrow i): Suppose that $U_\sigma \subseteq S$ for all $\sigma \in \mathcal{N}$. Note that Lemma 5.2 implies that U_σ is M -invariant, and we also have that the U_σ are mutually orthogonal since the sets B_σ are disjoint.

Suppose that $f \in S$. Then, since $\{B_\sigma\}_{\sigma \in \mathcal{N}}$ is a partition of Γ , it follows that $\widehat{f} = \sum_{\sigma \in \mathcal{N}} \widehat{f} \chi_{B_\sigma}$. This implies that $f \in \bigoplus_{\sigma \in \mathcal{N}} U_\sigma$ and consequently, S is the orthogonal direct sum

$$S = \bigoplus_{\sigma \in \mathcal{N}} U_\sigma.$$

As each U_σ is M -invariant, so is S . □

5.2. Characterization of M -invariance in terms of H -fibers.

In this section we will first express the conditions of Theorem 5.1 in terms of H -fibers. Then, we will give a useful characterization of the M -invariance for a finitely generated H -invariant space in terms of the Gramian.

If $f \in L^2(G)$ and $\sigma \in \mathcal{N}$, we define the function f^σ by

$$\widehat{f^\sigma} = \widehat{f} \chi_{B_\sigma}.$$

Let P_σ be the orthogonal projection onto S_σ , where

$$S_\sigma = \{f \in L^2(G) : \text{supp}(\widehat{f}) \subseteq B_\sigma\}.$$

Therefore

$$f^\sigma = P_\sigma f \quad \text{and} \quad U_\sigma = P_\sigma(S) = \{f^\sigma : f \in S\}.$$

Moreover, if $S = S_H(\mathcal{A})$ with \mathcal{A} a countable subset of $L^2(G)$, then

$$J_H(U_\sigma)(\omega) = \overline{\text{span}}\{\mathcal{T}_H(\varphi^\sigma)(\omega) : \varphi \in \mathcal{A}\}. \quad (9)$$

Remark 5.3. Note that the fibers

$$\mathcal{T}_H(\varphi^\sigma)(\omega) = \{\chi_{B_\sigma}(\omega + h^*)\widehat{\varphi}(\omega + h^*)\}_{h^* \in H^*}$$

can be described in a simple way as

$$\chi_{B_\sigma}(\omega + h^*)\widehat{\varphi}(\omega + h^*) = \begin{cases} \widehat{\varphi}(\omega + h^*) & \text{if } h^* \in \sigma + M^* \\ 0 & \text{otherwise.} \end{cases}$$

Therefore, if $\sigma \neq \sigma'$, $J_H(U_\sigma)(\omega)$ and $J_H(U_{\sigma'})(\omega)$ are orthogonal subspaces for a.e. $\omega \in \Omega$.

Combining Theorem 5.1 with Proposition 4.2 and (9) we obtain the following result.

Proposition 5.4. *Let S be an H -invariant space generated by a countable set $\mathcal{A} \subseteq L^2(G)$. The following statements are equivalent.*

- i) S is M -invariant.
- ii) $\mathcal{T}_H(\varphi^\sigma)(\omega) \in J_H(S)(\omega)$ a.e. $\omega \in \Omega$ for all $\varphi \in \mathcal{A}$ and $\sigma \in \mathcal{N}$.

Let $\Phi = \{\varphi_1, \dots, \varphi_\ell\}$ be a finite collection of functions in $L^2(G)$. Then, the Gramian G_Φ of Φ is the $\ell \times \ell$ matrix of H^* -periodic functions

$$\begin{aligned} [G_\Phi(\omega)]_{ij} &= \langle \mathcal{T}_H(\varphi_i)(\omega), \mathcal{T}_H(\varphi_j)(\omega) \rangle \\ &= m_{H^*}(\{0\}) \sum_{h^* \in H^*} \widehat{\varphi}_i(\omega + h^*) \overline{\widehat{\varphi}_j(\omega + h^*)} \end{aligned} \quad (10)$$

for $\omega \in \Omega$.

Given a subspace V of $L^2(G)$, the *dimension function* is defined by

$$\dim_V : \Omega \rightarrow \mathbb{N}_0 \cup \{\infty\}, \quad \dim_V(\omega) := \dim(J_H(V)(\omega)).$$

We will also need the next result which is a straightforward consequence of Proposition 4.1 and Proposition 4.2.

Proposition 5.5. *Let S_1 and S_2 be H -invariant spaces. If $S = S_1 \dot{\oplus} S_2$, then*

$$J_H(S)(\omega) = J_H(S_1)(\omega) \dot{\oplus} J_H(S_2)(\omega), \quad \text{a.e. } \omega \in \Omega.$$

The converse of this proposition is also true, but will not be needed.

Now we give a slightly simpler characterization of M -invariance for the finitely generated case.

Theorem 5.6. *If S is an H -invariant space, finitely generated by Φ , then the following statements are equivalent.*

- i) S is M -invariant.
- ii) For almost every $\omega \in \Omega$, $\dim_S(\omega) = \sum_{\sigma \in \mathcal{N}} \dim_{U_\sigma}(\omega)$.

iii) For almost every $\omega \in \Omega$, $\text{rank}[G_\Phi(\omega)] = \sum_{\sigma \in \mathcal{N}} \text{rank}[G_{\Phi^\sigma}(\omega)]$, where $\Phi^\sigma = \{\varphi^\sigma : \varphi \in \Phi\}$.

Proof. i) \Rightarrow ii): By Theorem 5.1, $S = \dot{\bigoplus}_{\sigma \in \mathcal{N}} U_\sigma$. Then, ii) follows from Proposition 5.5.

ii) \Rightarrow i): Since $\{B_\sigma\}_{\sigma \in \mathcal{N}}$ is a partition of Γ , $S \subseteq \dot{\bigoplus}_{\sigma \in \mathcal{N}} U_\sigma$. Then, by Remark 5.3 we have that

$$J_H(S)(\omega) \subseteq \dot{\bigoplus}_{\sigma \in \mathcal{N}} J_H(U_\sigma)(\omega).$$

Using ii), we obtain that $J_H(S)(\omega) = \dot{\bigoplus}_{\sigma \in \mathcal{N}} J_H(U_\sigma)(\omega)$. The proof follows as a consequence of Proposition 5.4.

The equivalence between ii) and iii) follows from (9). □

6. APPLICATIONS OF M -INVARIANCE

In this section we estimate the size of the supports of the Fourier transforms of the generators of a finitely generated H -invariant space which is also M -invariant.

We will not include the proofs of the results stated bellow, since they follow readily from the \mathbb{R}^d case (see [ACP09, Section 6]).

As a consequence of Theorem 5.6 we obtain the following theorem.

Theorem 6.1. *Let S be an H -invariant space, finitely generated by the set $\{\varphi_1, \dots, \varphi_\ell\}$, and define*

$$E_j = \{\omega \in \Omega : \dim_S(\omega) = j\}, \quad j = 0, \dots, \ell.$$

If S is M -invariant and \mathcal{D}' is any measurable section of Γ/M^ , then*

$$m_\Gamma(\{y \in \mathcal{D}' : \widehat{\varphi}_i(y) \neq 0\}) \leq \sum_{j=0}^{\ell} m_\Gamma(E_j) j \leq m_\Gamma(\Omega) \ell,$$

for each $i = 1, \dots, \ell$.

Corollary 6.2. *Let $\varphi \in L^2(G)$ be given. If $S_H(\varphi)$ is M -invariant for some closed subgroup M of G such that $H \not\subseteq M$, then $\widehat{\varphi}$ must vanish on a set of infinite m_Γ -measure.*

Since the uncertainty principle holds in LCA groups (see [Smi90]), we also can extend Proposition 6.4 of [ACP09] as follows.

Proposition 6.3. *If a nonzero function $\varphi \in L^2(G)$ has compact support, then $S_H(\varphi)$ is not M -invariant for any M closed subgroup of G such that $H \not\subseteq M$. In particular, if $x \in G$ and $x \notin H$,*

$$t_x \varphi \notin S_H(\varphi).$$

As a consequence of Theorem 6.1, in case that $M = G$, we obtain the following corollary.

Corollary 6.4. *If $\varphi \in L^2(G)$ and $S_H(\varphi)$ is G -invariant, then*

$$m_\Gamma(\text{supp}(\widehat{\varphi})) \leq m_\Gamma(\Omega).$$

6.1. Exactly M -invariance.

Let M be a closed subgroup of G containing a countable uniform lattice H . The next theorem states that there exists an M -invariant space S that is *not* invariant under any vector outside M . We will say in this case that S is *exactly* M -invariant.

Note that because of Proposition 3.1, an M -invariant space is exactly M -invariant if and only if it is not invariant under any closed subgroup M' containing M .

Theorem 6.5. *For each closed subgroup M of G containing a countable uniform lattice H , there exists a shift-invariant space of $L^2(G)$ which is exactly M -invariant.*

Proof. Suppose that $0 \in \mathcal{N}$ and take $\varphi \in L^2(G)$ satisfying $\text{supp}(\widehat{\varphi}) = B_0$, where B_0 is defined as in (4). Let $S = S(\varphi)$.

Then, $U_0 = S$ and $U_\sigma = \{0\}$ for $\sigma \in \mathcal{N}$, $\sigma \neq 0$. So, as a consequence of Theorem 5.1, it follows that S is M -invariant.

Now, if M' is a closed subgroup such that $M \subsetneq M'$, we will show that S can not be M' -invariant.

Since $M \subseteq M'$, $(M')^* \subseteq M^*$. Consider a section \mathcal{C} of the quotient $M^*/(M')^*$ containing the origin. Then, the set given by

$$\mathcal{N}' := \{\sigma + c : \sigma \in \mathcal{N}, c \in \mathcal{C}\},$$

is a section of $H^*/(M')^*$ and $0 \in \mathcal{N}'$.

If $\{B'_{\sigma'}\}_{\sigma' \in \mathcal{N}'}$ is the partition defined in (4) associated to M' , for each $\sigma \in \mathcal{N}$ it holds that $\{B'_{\sigma+c}\}_{c \in \mathcal{C}}$ is a partition of B_σ , since

$$B_\sigma = \Omega + \sigma + M^* = \bigcup_{c \in \mathcal{C}} \Omega + \sigma + c + (M')^* = \bigcup_{c \in \mathcal{C}} B'_{\sigma+c}. \quad (11)$$

We will show now that $U'_0 \not\subseteq S$, where U'_0 is the subspace defined in (5) for M' . Let $g \in L^2(G)$ such that $\widehat{g} = \widehat{\varphi} \chi_{B'_0}$. Then $g \in U'_0$. Moreover, since $\text{supp}(\widehat{\varphi}) = B_0$, by (11), $\widehat{g} \neq 0$.

Suppose that $g \in S$, then $\widehat{g} = \eta \widehat{\varphi}$ where η is an H^* -periodic function. Since $M \subsetneq M'$, there exists $c \in \mathcal{C}$ such that $c \neq 0$. By (11), \widehat{g} vanishes in B'_c . Then, the H^* -periodicity of η implies that $\eta(\gamma) = 0$ a.e. $\gamma \in \Gamma$. So $\widehat{g} = 0$, which is a contradiction.

This shows that $U'_0 \not\subseteq S$. Therefore, S is not M' -invariant. \square

REFERENCES

- [ACHKM09] A. Aldroubi, C. Cabrelli, C. Heil, K. Kornelson and U. Molter, *Invariance of a Shift-Invariant Space*, J. Fourier Anal. and Appl., to appear, 2009. (<http://arxiv.org/pdf/0707.2008>)
- [ACP09] M. Anastasio, C. Cabrelli and V. Paternostro, *Invariance of a Shift-Invariant Space in several variables*, preprint. (<http://arxiv.org/pdf/0905.1980>)

- [dBDR94a] C. de Boor, R. A. DeVore and A. Ron, *The structure of finitely generated shift-invariant spaces in $L^2(\mathbb{R}^d)$* , J. Funct. Anal. **119**, (1994), no. 1, 37-78.
- [dBDR94b] C. de Boor, R. A. DeVore and A. Ron, *Approximation from shift-invariant subspaces of $L^2(\mathbb{R}^d)$* , Trans. Amer. Math. Soc. **341**, (1994), 787-806.
- [Bow00] M. Bownik, *The Structure of Shift-Invariant Subspaces of $L^2(\mathbb{R}^d)$* , J. Funct. Anal. **177**, (2000), no. 2, 282-309.
- [CP09] C. Cabrelli and V. Paternostro, *Shift-Invariant Spaces on LCA Groups*, preprint. (<http://arxiv.org/pdf/0905.2373>)
- [FG64] J. Feldman and F. P. Greenleaf, *Existence of Borel Transversal in Groups*, Pacific J. Math. **25**, num. **3**, (1964).
- [Hel64] H. Helson, *Lectures on Invariant Subspaces*, Academic Press, New York-London, (1964).
- [HR63] Hewitt, E., Ross, K.A., *Abstract harmonic analysis Vol. I: Structure of topological groups Integration theory. Group representations*, Die Grundlehren der mathematischen Wissenschaften. 115. Berlin- Göttingen-Heidelberg: Springer-Verlag, VIII, 519 p. (1963).
- [HS64] M. Hasumi and T.P. Srinivasan, *Doubly invariant subspaces, II*, Pacific J. Math., **14**, (1964), 525-535.
- [KK98] E. Kanuith and G. Kutyniok, *Zeros or the Zak Transform on Locally Compact Abelian Groups*, American Mathematical Society **126**, num. **12**, (1998), 3561-3569.
- [KR08] R.A. Kamyabi Gol, R. Raisi Tousi, *The structure of shift invariant spaces on a locally compact abelian group* Journal of Mathematical Analysis and Applications, **340**, (2008), 219-225.
- [RS95] A. Ron and Z. Shen, *Frames and Stable Bases for Shift-Invariant Subspaces of $L^2(\mathbb{R}^d)$* , Canad. J. Math. **47**, (1995), no. 5, 1051-1094.
- [Rud62] W. Rudin, *Fourier Analysis on Groups*, John Wiley, (1962).
- [Smi90] K. T. Smith, *The uncertainty principle on groups*, SIAM J. Appl. Math., **50**, (1990), 876-882.
- [Sri64] T. P. Srinivasan, *Doubly invariant subspaces*, Pacific J. Math., **14**, (1964), 701-707.

(M. ANASTASIO) DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS EXACTAS Y NATURALES, UNIVERSIDAD DE BUENOS AIRES, CIUDAD UNIVERSITARIA, PABELLÓN I, 1428 BUENOS AIRES, ARGENTINA AND CONICET, CONSEJO NACIONAL DE INVESTIGACIONES CIENTÍFICAS Y TÉCNICAS, ARGENTINA

E-mail address: manastas@dm.uba.ar

(C. CABRELLI) DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS EXACTAS Y NATURALES, UNIVERSIDAD DE BUENOS AIRES, CIUDAD UNIVERSITARIA, PABELLÓN I, 1428 BUENOS AIRES, ARGENTINA AND CONICET, CONSEJO NACIONAL DE INVESTIGACIONES CIENTÍFICAS Y TÉCNICAS, ARGENTINA

E-mail address: cabrelli@dm.uba.ar

(V.PATERNOSTRO) DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS EXACTAS Y NATURALES, UNIVERSIDAD DE BUENOS AIRES, CIUDAD UNIVERSITARIA, PABELLÓN I, 1428 BUENOS AIRES, ARGENTINA AND CONICET, CONSEJO NACIONAL DE INVESTIGACIONES CIENTÍFICAS Y TÉCNICAS, ARGENTINA

E-mail address: vpater@dm.uba.ar